

scientific_name	range	type locality
<i>Smithornis capensis delacouri</i>	Sierra Leone to Ghana	Bandama, Ivory Coast.
<i>Smithornis capensis conjunctus</i>	Caprivi Strip and middle Zambezi Valley to nw Mozambique	30 miles west of the Victoria Falls, Zambesi R. (incomplete locality)
<i>Smithornis capensis capensis</i>	eastern South Africa (coastal KwaZulu-Natal and northeastern Eastern Cape)	Natal; restricted to the coastal forests of northern Zululand, Vincent, Bull. Brit. Orn. CL, 55, p. 96.
<i>Smithornis sharpei eurylaemus</i>	E Democratic Republic of the Congo	Moera, 40 kilometers northwest of Lake Albert Edward. 1
<i>Calyptomena viridis caudacuta</i>	S Myanmar, sw Thailand, Malay Peninsula and adjacent islands	India; error = Singapore Island, fide Chasen, 1935, p. 153.
<i>Calyptomena viridis siberu</i>	Mentawai Is. (Siberut, North Pagai and South Pagai)	Siberut Island.
<i>Psarisomus dalhousiae cyanicauda</i>	Southeast Thailand and Cambodia	Kao Sabab, southeastern Siam.
<i>Psarisomus dalhousiae divinus</i>	S Vietnam (s Annam)	Lang Bian Mts., Haut-Donai Province, Annam.
<i>Psarisomus dalhousiae psittacinus</i>	Malay Peninsula and Sumatra	Sumatra.
<i>Corydon sumatranus laoensis</i>	S Myanmar to n Thailand and Indochina	the Siamese Lao; type from Doi Souteb, 3300 feet.
<i>Corydon sumatranus sumatranus</i>	Extreme s Thailand, Malay Peninsula, Sumatra and Penang I.	interior of Sumatra.
<i>Corydon sumatranus brunnescens</i>	Northwest Borneo (Sarawak) and North Natuna Islands	Baram, Borneo.
<i>Serilophus lunatus elisabethae</i>	SE China (se Yunnan and sw Guangxi) and e Tonkin	Hokow, S.E. Yunnan (alt. 300 ft.)
<i>Serilophus lunatus polionotus</i>	Mountains of Hainan I.	Mt. Wuchi, Hainan.
<i>Serilophus lunatus stolidus</i>	S Myanmar and peninsular Thailand (except extreme south)	Kao Nawng, 1200-1500 feet, [= Khao Nong, a mountain at lat. 8° 55' N., long. 99° 38' E.], Bandon, peninsular Siam.
<i>Serilophus lunatus impavidus</i>	S Laos (Bolaven Plateau)	Thateng, lat. 15° 31' N., long. 106° 22' E., Saravane Province, Bas-Laos.
<i>Serilophus lunatus rothschildi</i>	Malay Peninsula and extreme s peninsular Thailand	Gunong Ijau, 3000 feet, Perak.
<i>Serilophus lunatus intensus</i>	Sumatra	Siolak Dras, 3100 feet, Korinchi [Kerintji] Valley, Sumatra.
<i>Cymbirhynchus macrorhynchus affinis</i>	Southwest Myanmar	Arakan.
<i>Cymbirhynchus macrorhynchus malaccensis</i>	Extreme s Thailand and Malay Peninsula	Malacca.
<i>Eurylaimus javanicus friedmanni</i>	SE Myanmar, Thailand and Indochina	Sathani Hin Lap, lat. 14° 40' N., long. 101° 10' E., Siam.
<i>Eurylaimus javanicus pallidus</i>	S Thailand (Isthmus of Kra) and Malay Peninsula	Kao Nawng [= Khao Nong, a mountain at lat. 8° 55' N., long. 99° 38' E.], Peninsular Siam.
<i>Eurylaimus javanicus harterti</i>	Sumatra, Riau Archipelago, Bangka I. and Belitung I.	Deli, Sumatra.
<i>Eurylaimus javanicus javanicus</i>	Java	Java.
<i>Erythropitta erythrogaster yairocho</i>	Sulu Islands, Philippines	Sibutu, Sulu Islands.
<i>Erythropitta erythrogaster inspeculata</i>	Talaut Islands	Talaut (Kabruang and Salibabu).
<i>Erythropitta caeruleitorques</i>	Sangihe I. (ne of Sulawesi)	Petta, Sangir Islands.
<i>Erythropitta palliceus</i>	Siau and Tahulandang islands (Celebes Sea)	Siau, Sangir Islands.
<i>Erythropitta celebensis</i>	Sulawesi, Manterawu I. and Togian Islands	Tondano, Celebes.
<i>Erythropitta dohertyi</i>	Banggai Is. (Peleng, Banggai) and Sula Is. (Taliabu, Seho, Mangole)	Sula Mangoli.
<i>Erythropitta rufiventris rufiventris</i>	Morotai, Halmahera, Kasiruta, Bacan, Moti, Damar and Obi islands	unknown; Batjan designated by Hartert, 1903, Novit. Zool., 10, pp. 13, 57.
<i>Erythropitta rufiventris cyanonota</i>	Ternate I. (North Moluccas)	Ternate.
<i>Erythropitta rubrinucha rubrinucha</i>	Buru I. (South Moluccas)	Buru.
<i>Erythropitta rubrinucha piroensis</i>	Seram I. (South Moluccas)	Piru, Ceram.

<i>Erythropitta macklotii macklotii</i>	Aru Islands, Raja Ampat Islands, and western and southern New Guinea	Lobo, Triton Bay, southwestern New Guinea.
<i>Erythropitta macklotii finschii</i>	D'Entrecasteaux Archipelago (Fergusson and Goodenough Islands; recent sight records from Normanby Island probably also refer to this species), off of southeastern New Guinea	Astrolabe Range, New Guinea; error for Fergusson Island, D'Entrecasteaux Archipelago, fide Hartert, 1895, Novit. Zool., 2, p. 61.
<i>Erythropitta novaehibernicae extima</i>	New Hanover (Bismarck Archipelago)	New Hanover, Bismarck Archipelago.
<i>Erythropitta splendida</i>	Tabar I. (e of New Ireland)	Tabar Island, Tabar group.
<i>Erythropitta gazellae</i>	S Bismarck Archipelago (New Britain and adjacent islands)	Ralum, New Britain.
<i>Erythropitta meeki</i>	Rosell I. (Louisiade Archipelago)	Rosell Island.
<i>Erythropitta granatina coccinea</i>	Malay Peninsula, s peninsular Thailand, s Myanmar and Sumatra	Malacca.
<i>Hydrornis oatesi oatesi</i>	Myanmar to ne Laos and sw Thailand	Toungoo district, Upper Pegu.
<i>Hydrornis oatesi deborah</i>	southern Thai-Malay Peninsula	Fraser's Hill, 1,200 m., 3° 43' N, 101° 44' E, Ulu Tranum Forest Reserve, Main Range, Pahang.
<i>Hydrornis nipalensis hendeei</i>	S China (se Yunnan, sw Guangxi) to n Vietnam and n Laos	Muong-moun, Tonkin.
<i>Hydrornis soror flynnstonei</i>	SE Thailand and sw Cambodia	Khao Soi Dao Tai, 4800 ft, Pong Nam Ron district, Chanthaburi Province, south-eastern Thailand, coordinates 12:56N, 102:12E
<i>Hydrornis caeruleus caeruleus</i>	S Myanmar, s Thailand, Malay Peninsula and Sumatra	Sumatra.
<i>Hydrornis schneideri</i>	Batak Mountains of n Sumatra (critically endangered)	Batak Mountains, upper Deli, North Sumatra.
<i>Hydrornis schwaneri</i>	Borneo	Borneo.
<i>Hydrornis cyaneus cyaneus</i>	E Bangladesh, ne India, s China, Myanmar, Thailand, Indochina	Arracan.
<i>Hydrornis cyaneus willoughbyi</i>	Mts. of c Laos to s Annam	Xiengkhouang, Laos.
<i>Pitta sordida abbotti</i>	Nicobar Islands	Great Nicobar.
<i>Pitta sordida bangkana</i>	Bangka and Belitung islands (off Sumatra)	Bangka Island.
<i>Pitta sordida sanghirana</i>	Sangihe I. (ne of Sulawesi)	Sangir.
<i>Pitta sordida forsteni</i>	northern Sulawesi (Minahasa Peninsula)	Tondano, Celebes. (Not Citta Melanocephal. [a] Wagler, 1827.)
<i>Pitta sordida goodfellowi</i>	Aru Islands (off s New Guinea)	Silbattabatta, Aru Islands.
<i>Pitta sordida mefoorana</i>	Numfor I. (off nw New Guinea)	Mefoor Island.
<i>Pitta sordida rosenbergii</i>	Biak I. (off nw New Guinea)	Soek Island.
<i>Pitta steerii steerii</i>	S Philippines (Mindanao)	Dumalon, Mindanao.
<i>Pitta concinna</i>	Lombok, Sumbawa, Flores, Adonara, Lombok and Alor islands	Lombok
<i>Pitta elegans maria</i>	Lesser Sundas (Sumba and s Flores)	Sumba Island.
<i>Pitta elegans virginalis</i>	Tanahjampea, Kalaotoa and Kalao islands (off Sulawesi)	Djampea Island.
<i>Pitta elegans elegans</i>	Sangihe I., Sula Islands and c Moluccas	Timor.
<i>Pitta vigorsii</i>	Seram, Watubela, Banda, Tayandu, Kai and Tanimbar islands	eastern coast of Australia; error, corrected to Banda by Wallace, 1862, Proc. Zool. Soc. London, p. 188
<i>Pitta anerythra pallida</i>	Bougainville (n Solomon Islands)	Bougainville.
<i>Pitta iris johnstoneiana</i>	NW Western Australia (nw Kimberley region)	Mt Trafalgar, Prince Regent River Reserve, WA
<i>Euchrepomis callinota venezuelana</i>	NW Venezuela (Sierra de Perijá), Mérida and Barinas	Cerro Pejochaina, Sierra de Perijá, Zulia, Venezuela.
<i>Batara cinerea excubitor</i>	E slope of Andes of c Bolivia (w Santa Cruz)	Samaipata, Santa Cruz, Bolivia.

<i>Sakesphorus canadensis canadensis</i>	Suriname and coastal French Guiana	Canada; error = Cayenne, by designation of Berlepsch and Hartert, Nov. Zool., 9, 1902, p. 70.
<i>Thamnophilus doliatus cadwaladeri</i>	S Bolivia (Tarija)	Villa Montes, 2000 feet, Tarija, Bolivia.
<i>Thamnophilus multistriatus selvae</i>	W slope of Western Andes of Colombia (upper Río San Juan)	La Selva, 6000 feet, Caldas, Colombia.
<i>Thamnophilus multistriatus multistriatus</i>	Colombia (e slope of Cent. Andes and w slope of E and S Andes)	Colombia = Bogota.
<i>Thamnophilus multistriatus oecotonophilus</i>	W slope of E Andes of Colombia and Sierra de Perijá (w Venezuela)	Soatá, Departamento de Boyacá, Colombia; alt. 1700-2040 m
<i>Thamnophilus tenuepunctatus tenuepunctatus</i>	E slope of Eastern Andes of n-cent. Colombia	Anolaima, Colombia.
<i>Thamnophilus atrinucha gorgonae</i>	Gorgona I. (off w Colombia)	Gorgona Island.
<i>Thamnophilus insignis nigrofrontalis</i>	Extreme sw Venezuela (Macizo de Sipapo in w Amazonas)	Cerro Paraque, 1600 meters, Territorio Amazonas, Venezuela.
<i>Thamnistes anabatinus gularis</i>	Extreme nw Venezuela (Táchira) and (? adj. ne Colombia)	Cerro El Tetéo, Burgua, Estado Táchira, Venezuela; 1250 meters
<i>Epinecrophylla haematonota ssp. nov.</i>	N of Río Marañón, W of Río Napo	N/A
<i>Epinecrophylla ornata ornata</i>	Andean foothills of central Colombia (Meta)	Nova Granada i.e. Bogota, Colombia.
<i>Epinecrophylla ornata saturata</i>	S-cent. Colombia to e Ecuador and ne Peru (n of R. Marañón)	upper Río Suno, eastern Ecuador.
<i>Myrmotherula longicauda soderstromi</i>	E slope of Andes of s Colombia and n Ecuador (Napo)	near Río Napo, eastern Ecuador.
<i>Myrmotherula longicauda longicauda</i>	Central Peru (Junín)	Chontabamba, La Merced, and La Gloria, Peru.
<i>Myrmotherula sunensis yessupi</i>	E-central Peru (Huánuco, Pasco) and sw Amazonian Brazil	Puerto Yessup, Junin, Peru.
<i>Myrmotherula behni camanii</i>	S Venezuela (Cerro Camani in n Amazonas)	Cerro Camani, Territorio Amazonas, Venezuela, 1250 meters
<i>Herpsilochmus roraimae kathleenae</i>	Tepuis of sw Venezuela and adj. Brazil (n Amazonas)	Caño Laguna, Falda occidental del Cerro Parú, Amazonas Venezuela.
<i>Terenura sicki</i>	E Brazil (e Alagoas and e Pernambuco)	Sierra Blanca, 9°15'S, 35°50'W, Murici, Alagoas, Noreste del Brasil.
<i>Gymnocichla nudiceps sanctamartae</i>	N Colombia (Santa Marta region and Magdalena Valley)	Santa Marta, Colombia.
<i>Sclateria naevia diaphora</i>	S-cent. Venezuela (lower R. Caura drainage in nw Bolívar)	Río Mocho, Río Caura, Venezuela.
<i>Myrmeciza longipes boucardi</i>	Colombia (upper Magdalena Valley from Cundinamarca to Huila)	Bogota.
<i>Poliocrania exsul maculifer</i>	Pacific slope of w Colombia and w Ecuador	Paramba, 3500 feet, Esmeraldas, Ecuador.
<i>Myrmophylax atrothorax metae</i>	Central Colombia (Meta, w Guaviare)	Quenane, Llanos de Meta, Colombia.
<i>Myrmophylax atrothorax maynana</i>	N-cent. Peru (south of Río Marañón and west of Río Huallaga)	Yurimaguas, Peru.
<i>Gymnopithys bicolor daguae</i>	Pacific slope of w Colombia (s Chocó to Cauca)	El Paillon [= Pailon], near Buenaventura, Valle, Colombia.
<i>Gymnopithys bicolor ruficeps</i>	Tropical central Colombia (Antioquia)	Cauca Valley = Río Nechi, Antioquia, Colombia.
<i>Gymnopithys rufigula pallidigula</i>	Tepuis of extreme s Venezuela (Pica Yavita-Pimichín environs)	Campamento la Cruz, 150 meters, Yavita-Pimichin portage, Amazonas, Venezuela.
<i>Pittasoma michleri zeledoni</i>	Caribbean slope of Costa Rica and w Panama	Río Sucio, Costa Rica.
<i>Pittasoma rufopileatum rosenbergi</i>	W Andes of Colombia and Baudó Mountains (south to Chocó)	Sipi, 150 feet, Río Sipi, Choco, Colombia.
<i>Pittasoma rufopileatum harterti</i>	Lowlands and foothills of Colombia (Cauca and w Nariño)	Barbacoas, Nariño, Colombia.
<i>Conopophaga castaneiceps chocoensis</i>	Colombia (w slope of W Andes and Baudó Mts. in w Chocó)	Baudó Mts., 3500 feet, Choco, Colombia.
<i>Grallaria gigantea lehmanni</i>	Andes of sw Colombia (head of Magdalena Valley)	San Marcos, 3000 meters, Moscopan, Cauca, Colombia.

<i>Grallaria gigantea hyalodroma</i>	SW Colombia (Nariño) and W slope of Andes of Ecuador	Gualea, 6000 feet, Pichincha, Ecuador.
<i>Grallaria excelsa phelpsi</i>	Coastal mts. n Venezuela (Colonia Tovar in Aragua)	Colonia Tovar, 5900 feet, Aragua, Venezuela.
<i>Grallaria varia intercedens</i>	E Brazil (Pernambuco and s Bahia to se Minas Gerais)	Baia; type a trade skin.
<i>Grallaria alleni andaquiensis</i>	Andes of Colombia (Magdalena) to Ecuador (Napo, Cotopaxi)	Inmediaciones de la Cueva de los Guácharos, valle del Río Suaza, Acevedo, Huila, Colombia.
<i>Grallaria guatimalensis aripoensis</i>	Trinidad; popul	Aripo Mts., 2000 feet, Trinidad.
<i>Grallaria guatimalensis carmelitae</i>	Santa Marta Mts. (ne Colombia) and Sierra de Perijá	Pueblo Viejo, Santa Marta, Colombia.
<i>Grallaria haplonota pariae</i>	NE Venezuela (subtropical mts. of Paría Peninsula)	Cerro Azul, Cristobal Colon, Paria Peninsula, Sucre, Venezuela.
<i>Grallaria ruficapilla perijana</i>	Sierra de Perijá (Colombia/Venezuela border)	La Sabana, 1500 meters, Rio Negro, Perija District, Zulia, Venezuela.
<i>Grallaria ruficapilla nigrolineata</i>	Andes of w Venezuela (Táchira, Mérida and Trujillo)	Merida, Venezuela.
<i>Grallaria ruficapilla connectens</i>	Subtropical Andes of sw Ecuador	Taraguacocha, 9750 feet, Cordillera de Chilla, El Oro, Ecuador.
<i>Grallaria griseonucha griseonucha</i>	Andes of w Venezuela (e Mérida)	Paramo de Culata, Andes of Merida, Venezuela.
<i>Grallaria nuchalis ruficeps</i>	Colombia (Central Andes and w slope of Eastern Andes)	Antioquia, Colombia; type from Medellin.
<i>Grallaria nuchalis obsoleta</i>	W slope of Andes of nw Ecuador (Imbabura and Pichincha)	west side of Mt. Pichincha, 12,000 feet, Ecuador.
<i>Grallaria hypoleuca hypoleuca</i>	W slope of Eastern Andes of Colombia	Santa Fe de Bogota.
<i>Grallaria spatiator</i>	Santa Marta Mountains (ne Colombia)	Macotama, 8000 feet, Sierra Nevada de Santa Marta, Colombia
<i>Grallaria rufocinerea romeroana</i>	S Colombia to Andes of ne Ecuador (nw Sucumbíos)	Inmediaciones de la Cascada del Río Bedón (o Río San José), margen derecha, Parque Nacional Natural del Puracé, 76°17'W, 2°20' N., La Plata, Huila, Colombia.
<i>Grallaria quitensis atuensis</i>	Andes of n Peru (s Amazonas and e La Libertad)	Atuen, 12,000 feet, Amazonas, Peru.
<i>Grallaria milleri gilesi</i>	northern Central Andes of Colombia (Antioquia); known from a single 19th century specimen, probably extinct	Santa Elena, 8 km. al este de Medellín, 06 15'N, 75 35'W., Antioquia, Colombia.
<i>Hylopezus perspicillatus pallidior</i>	Colombia (lower Cauca, upper Sinú and Magdalena valleys)	El Tambor, Santander, Colombia.
<i>Hylopezus perspicillatus lizanoi</i>	Pacific slope of sw Costa Rica	Trojas, Costa Rica.
<i>Myrmothera fulviventris caquetae</i>	Lowlands of extreme se Colombia (w Caquetá)	Morelia, Caqueta, Colombia
<i>Myrmothera campanisona modesta</i>	Base of Eastern Andes of se Colombia (south from Meta)	Santa Fe de Bogota.
<i>Grallaricula flavirostris costaricensis</i>	Mountains of Costa Rica and w Panama (east to Veraguas)	Barranca, Costa Rica.
<i>Grallaricula cucullata cucullata</i>	Central and Eastern Andes of Colombia	Bogota.
<i>Grallaricula cucullata venezuelana</i>	Andes of central Colombia and nw Venezuela	Río Chiquito, Hacienda La Providencia, Táchira, Venezuela.
<i>Grallaricula ferruginepectus rara</i>	eastern Andes of Colombia, and the Sierra de Perijá (Colombia/ Venezuela border)	Llanos de Medina, Cundinamarca, Colombia.
<i>Grallaricula nana olivascens</i>	coastal mountains of north central Venezuela (Aragua and Distrito Federal)	Galipan, 2000 meters, Cerro del Avila, near Caracas, Venezuela.
<i>Grallaricula nana nana</i>	Eastern Andes of Colombia	Colombia = Bogota.
<i>Grallaricula cumanensis pariae</i>	NE Venezuela (subtropical mountains of Paría Peninsula)	Cerro Azul, Cristobal Colon, Paria Peninsula, Sucre, Venezuela.
<i>Scelorchilus albicollis atacamae</i>	N Chile (sw Antofogasta and Atacama to n Coquimbo)	Caldera, Atacama, Chile.
<i>Scelorchilus rubecula mochae</i>	Mocha I. (off Chile)	Mocha Island, Chile.
<i>Scytalopus atratus confusus</i>	Cent. Andes and e slope of W Andes of Colombia	Miraflores, 6800 feet, east of Palmira, Colombia.

<i>Scytalopus rodriguezi yariguiorum</i>	northern Colombia (Serranía de los Yariguíes, Santander)	Camino del Lenguerke' above Finca El Taliman, Vereda El Centro, San Vicente de Chucuri municipality, west slope of the Yariguíes massif, dpto. Santander, Colombia (06°50'N, 73°2'W; 2,100 m). Serranía de los Yariguíes, Colombia
<i>Scytalopus alvarezlopezi</i>	Pacific slope of Colombian Andes (western Antioquia south to southwestern Valle del Cauca)	Colombia, department of Risaralda, municipality of Mistrato, Alto de Pisones, 8 km northeast of Geguadas, elevation 1,710 m, shallow ravine in cloud forest; coordinates 5 25 N, 76 02 W.
<i>Scytalopus griseicollis gilesi</i>	Andes of northern Colombia (Serranía de los Yariguíes, Santander)	Páramo Filo Pamplona, arriba de La Aurora, vereda San Isidro, 06 38'N; 73 24'W, Municipio de Galán, Santander, Colombia.
<i>Formicarius analis virescens</i>	West base of Santa Marta Mts. (ne Colombia)	Fundacion, Santa Marta, Colombia.
<i>Formicarius analis griseoventris</i>	Mts. of n Colombia and nw Venezuela in w Maracaibo basin	Kunana, Sierra de Perijá, Zulía, Venezuela
<i>Formicarius analis connectens</i>	E Colombia east of the Andes	Villavicencio, 1600 feet, Colombia.
<i>Formicarius nigricapillus nigricapillus</i>	Caribbean slope of e Costa Rica and both slopes of Panama	Buena Vista, Costa Rica.
<i>Formicarius rufipectus carrikeri</i>	W and Central Andes of Colombia and w Ecuador	San Antonio, 6600 feet, Western Andes, Colombia.
<i>Chamaeza campanisona yavii</i>	Tepuis of south-central Venezuela (Cerro Yaví)	Cerro Yaví, 1600 meters, Territorio Amazonas, Venezuela.
<i>Chamaeza campanisona huachamacarii</i>	Tepuis of s Venezuela (Cerro Huachamacari)	Cerro Huachamacare, Amazonas, Venezuela.
<i>Chamaeza campanisona obscura</i>	Subtropical mountains of se Venezuela (Bolívar and s Amazonas)	Mt. Auyan-tepui, 1100 meters, Bolívar, Venezuela.
<i>Sclerurus obscurior bahiae</i>	E Brazil (Alagoas to ne São Paulo)	Baía, Brazil
<i>Sclerurus guatemalensis ennosiphyllus</i>	Colombia (Río Magdalena Valley in Santander)	Volador, cerca de El Tigre, 25 millas al oeste de Simití, Bolívar, Colombia.
<i>Sclerurus caudacutus umbretta</i>	Coastal e Brazil (Alagoas to Espírito Santo)	Baía.
<i>Sclerurus caudacutus calligineus</i>	Alagoas and Pernambuco	BRAZIL: Alagoas; Usina Mangabeiras
<i>Sclerurus albigularis canigularis</i>	Montane forests of Costa Rica and w Panama (Chiriquí)	Turrialba, Costa Rica.
<i>Sclerurus albigularis kunanensis</i>	NE Venezuela (Paría Peninsula)	Kunana, Sierra de Perijá, Río Negro, Venezuela.
<i>Geositta cunicularia fissirostris</i>	C Chile (s Atacama to Llanquihué)	Valparaiso, Chile.
<i>Geositta cunicularia contrerasi</i>	W-c Argentina (Sierra Grandes in Córdoba)	Los Gigantes, Sierras Grandes de Córdoba, Córdoba, Argentina.
<i>Geositta cunicularia hellmayri</i>	W Argentina	Huanuluan, Río Negro, Argentina.
<i>Geositta rufipennis harrisoni</i>	N Chile (sw Antofagasta)	Quebrada Paposo; 22° 02' S, 70° 29'W, Provincia de Antofagasta, Chile.
<i>Geositta rufipennis ottowi</i>	W-c Argentina (Sierra de Córdoba)	Sierra de Córdoba, Córdoba, Argentina.
<i>Geositta rufipennis hoyi</i>	W Argentina (Mendoza to Neuquén) and s Chile (s Aysén)	Quebrada de las Lajas, Departamento Las Heras, Mendoza, Argentina.
<i>Geositta rufipennis gaii</i>	Southwest Argentina (s Neuquén to Chubut)	Cerro Carbón, Valle del Arroyo Bernal, 9 Km. del Lago Nahuel Huapi, Bariloche, Río Negro, Argentina.
<i>Deconychura longicauda minor</i>	Tropical n Colombia	El Tambor, Santander, Colombia.
<i>Dendrocincla tyrannina hellmayri</i>	Andes of e Colombia and w Venezuela (sw Táchira)	Paramo de Tama, Colombia.
<i>Dendrocincla fuliginosa lafresnayeii</i>	N and e Colombia and adjacent nw Venezuela	Upper Amazon? error = Santa Marta region by designation of Chapman, Bull. Am. Mus. Nat. Hist., 36, 1917, p. 418.
<i>Glyphorhynchus spirurus amacurensis</i>	NE Venezuela (Delta Amacuro)	Jobure, Territorio Delta Amacuro, Venezuela.
<i>Dendrocolaptes sanctithomae sheffleri</i>	Pacific slope of sw Mexico (Guerrero and Oaxaca)	19 mi. N Puerto Angel, Oaxaca, México, elevation 900 feet

<i>Dendrocolaptes sanctithomae punctipectus</i>	N Colombia and nw Venezuela (Zulia and n Mérida)	La Sierra, Rio Negro, Perija' district, Zulia, Venezuela.
<i>Dendrocolaptes picumnus puncticollis</i>	Highlands of s Mexico (Chiapas) to Guatemala and w Honduras	Tactic and San Geronimo, Vera Paz, Guatemala.
<i>Dendrocolaptes picumnus casaresi</i>	Andean foothills of nw Argentina (Jujuy, Salta and Tucumán)	Concepcion, Tucuman.
<i>Xiphocolaptes promeropirhynchus omiltemensis</i>	Subtropical sw Mexico (Sierra Madre del Sur of Guerrero)	Omilteme, Guerrero, Mexico.
<i>Xiphocolaptes promeropirhynchus costaricensis</i>	Highlands of c Costa Rica and sw Panama (Chiriquí)	Naranjo, Cartago, Costa Rica.
<i>Xiphocolaptes promeropirhynchus panamensis</i>	Mountains of Pacific slope of s Panama (Veraguas)	Chitra, 3600 feet, Veraguas, Panama.
<i>Xiphocolaptes promeropirhynchus sanctaemartae</i>	N Colombia (Santa Marta Mts.)	San Lorenzo, Santa Marta Mountains, Colombia.
<i>Xiphocolaptes promeropirhynchus virgatus</i>	W slope of C Andes of Colombia east to Río Magdalena	locality unknown; presumably from Antioquia, Colombia.
<i>Xiphocolaptes promeropirhynchus procerus</i>	Mountains of n and c Venezuela	Caracas.
<i>Xiphocolaptes promeropirhynchus macarenae</i>	W-c Colombia (Macarena Mts. and foothills of Cent. Andes)	Ladera este del Pico Rengifo, La Macarena, Meta, Colombia.
<i>Xiphocolaptes falcirostris franciscanus</i>	E Brazil (w Bahia and n and nw Minas Gerais)	Brejo Januaria, left bank of Rio Sao Francisco, Minas Gerais.
<i>Xiphocolaptes albicollis villanovae</i>	Known only from type locality in ne Brazil (ne Bahia)	Vila Nova da Raihna (actual Bonfim) en el alto Río Itapicuru, Bahía, Brasil.
<i>Xiphocolaptes albicollis bahiae</i>	E Brazil (e and c Bahia)	Macaco Secco, near Andarahy, Baia.
<i>Xiphorhynchus obsoletus caicarae</i>	Middle Orinoco Valley in cent. Venezuela (nw Bolívar)	Caicara, 07°37'S, 66°10'W, margen derecha del bajo Río Orinoco, Bolívar, Venezuela.
<i>Xiphorhynchus elegans buenavistae</i>	E slope of E Andes of Colombia in upper Orinoco drainage	Buena Vista, 04°10'S, 73°41'W, cerca de Villavicencio, Meta, Colombia
<i>Xiphorhynchus susurrans rosenbergi</i>	W Colombia (upper Cauca Valley in Valle)	Guabinas, Rio Cauca, Colombia.
<i>Xiphorhynchus susurrans margaritae</i>	Isla Margarita (off n Venezuela)	El Cafetal, 740 meters, Margarita Island, Venezuela.
<i>Xiphorhynchus guttatus gracilirostris</i>	ne Brazil (Serra de Baturité, Ceará)	Sierra de Baturité, Ceará, Brasil.
<i>Xiphorhynchus lachrymosus alarum</i>	Tropical n Colombia (Sinú, Cauca and Magdalena valleys)	Puerto Valdivia, 360 feet, lower Cauca Valley, Colombia.
<i>Xiphorhynchus triangularis hylodromus</i>	Coastal and interior mts. of n Venezuela	Cordillera de la Costa, 4500 feet, above Rancho Grande, Aragua, Venezuela.
<i>Dendroplex picus dugandi</i>	NW Colombia (Chocó, Bolívar, Atlántico and Santa Marta Mts.)	Puerto Sagoc, La Gloria, Magdalena, Colombia.
<i>Dendroplex picus paraguanae</i>	NW Venezuela (Falcón, n Lara)	Adícora, Península de Paraguaná, Falcón, Venezuela.
<i>Dendroplex picus choicus</i>	N Venezuela (e Falcón east to Miranda)	Borburata, Puerto Cabello, Carabobo, Venezuela.
<i>Dendroplex picus saturator</i>	Tropical e Colombia to w Venezuela (cent. and s Maracaibo basin)	Encontrados, Zulia, Venezuela.
<i>Dendroplex picus altirostris</i>	Trinidad	Trinidad.
<i>Campylorhamphus pusillus tachirensis</i>	Andes of extreme ne Colombia and nw Venezuela (Zulia, Táchira)	Hacienda la Providencia, Río Chiquito, Táchira, Venezuela.
<i>Campylorhamphus pusillus guapiensis</i>	Coastal lowlands of sw Colombia (Cauca)	San Antonio de Guajui, 2°34'S, 77°54'W., bajo Río Guapi, Cauca, Colombia.
<i>Lepidocolaptes souleyetii guerrerensis</i>	W Mexico (Sierra Madre del Sur of Guerrero and sw Oaxaca)	Rincon, Guerrero, Mexico.
<i>Lepidocolaptes souleyetii uaireni</i>	Extreme se Venezuela (along Rio Uairén in se Bolívar)	Hato Santa Teresa, Rio Uairen, 940 meters, Bolivar, Venezuela.
<i>Lepidocolaptes lacrymiger sanctaemartae</i>	Santa Marta Mts. of n Colombia	Valparaiso, 5000 feet, Sierra Nevada de Santa Marta, Colombia.
<i>Lepidocolaptes lacrymiger lafresnayi</i>	Coastal cordillera of n Venezuela	Caracas.

<i>Lepidocolaptes lacrymiger sneiderni</i>	Andes of Colombia	El Tambo, 6000 feet, east slope of the Western Andes, Cauca, Colombia.
<i>Lepidocolaptes lacrymiger frigidus</i>	E slope of Andes of s Colombia (Nariño)	Cordillera de Guamués, Nariño, Colombia.
<i>Lepidocolaptes lacrymiger aequatorialis</i>	Pacific slope of Andes of sw Colombia (Nariño) and Ecuador	Banos, Ecuador.
<i>Xenops minutus olivaceus</i>	Lowlands of ne Colombia	Kunana, 1100 m., Rio Negro, Zulia, Venezuela
<i>Xenops rutilans septentrionalis</i>	Highlands of Costa Rica and w Panama	Guayabo, Costa Rica.
<i>Xenops rutilans incomptus</i>	E Panama (Darién)	Cana, Cerro Pirre, Darién, Panamá
<i>Xenops rutilans phelpsi</i>	Santa Marta Mountains (ne Colombia)	Las Vegas, Santa Marta, Colombia.
<i>Ochetorhynchus melanurus atacamae</i>	Mountains of n-c Chile (Atacama and Coquimbo)	Domeyko, 63 kilom. south of Vallenar, Atacama, Chile.
<i>Tarphonimus certhioides luscini</i>	Chaco of w Argentina (La Rioja and Mendoza)	middle and western region of Argentina; types from Mendoza.
<i>Furnarius leucopus endoecus</i>	N Colombia (Magdalena Valley) and w Venezuela (s Zulia)	Encontrados, Zulia, Venezuela.
<i>Lochmias nematura nelsoni</i>	Extreme e Panama (Darién)	Mt. Pirri, 4500 feet, near head of Rio Limon, eastern Panama.
<i>Phleocryptes melanops loaensis</i>	N Chile (Tarapacá)	Chacance, junction of the Rio San Salvador with the Rio Loa, lat. 22° 24' S., long. 69° 32' W., 1204 meters, Dept. Tocopilla, Chile.
<i>Upucerthia dumetaria peruana</i>	Andes of s Peru (Puno)	15 millas al este de Juliaca, Puno, Perú.
<i>Cinclodes pabsti espinhacensis</i>	east central Brazil (Serra do Espinhaço Meridional, Minas Gerais)	Tainhas, Rio Grande do Sul, Espinhaço Range, southeastern Brazil
<i>Cinclodes albidiventris oreobates</i>	Santa Marta Mts., C and Eastern Andes of n Colombia	Sierra Nevada de Santa Marta, Colombia.
<i>Cinclodes albidiventris heterurus</i>	Andes of w Venezuela (Mérida, Trujillo and s Lara)	Culata, San Antonio, and Valle, Venezuela.
<i>Cinclodes albiventris rufus</i>	Andes of nw Argentina (Campo de Arenal in Catamarca)	Angostura-Hualfin, Belém, Catamarca, Argentina.
<i>Cinclodes oustaleti baekstroemii</i>	Juan Fernández Islands (off Chile)	Islands of Mas Afuera and Mas a Tierra.
<i>Cinclodes oustaleti hornensis</i>	Tierra del Fuego and Cape Horn Archipelago	Hermit Island, west of Cape Horn.
<i>Cinclodes excelsior columbianus</i>	N Colombia (Santa Marta Mountains and Nevado de Tolima)	Paramo de Santa Isabel, 12,700 feet, Central Andes, Colombia.
<i>Cichlocolaptes mazarbarnetti</i>	northeastern Brazil; currently known from only two sites, in Alagoas and Pernambuco	Serra Branca, Murici, 09° 15' S, 35° 50' W, 550 m above sea level, Alagoas State, Brazil
<i>Philydor novaesi</i>	Montane forests of ne Brazil (Alagoas)	Sierra Blanca, Município de Murici, Alagoas, Brasil.
<i>Syndactyla dimidiata dimidiata</i>	SW Brazil (s Mato Grosso)	Sangrador and Rio Manso, Mato Grosso.
<i>Syndactyla roraimae paraquensis</i>	tepuis of southern Venezuela (Cerro Sipapo [Cerro Paraque], northwestern Amazonas)	Cerro Paraque, 1450 meters, Territorio Amazonas, Venezuela.
<i>Dendroma rufa cuchiverus</i>	Subtropical s Venezuela (Bolívar)	Cerro El Negro, upper Rio Cuchivero, 1000 meters, Bolivar, Venezuela
<i>Clibanornis rubiginosus fumosus</i>	Subtropical w Panama (Chiriquí)	Bibalaz, Volcan de Chiriqui, Panama.
<i>Clibanornis rubiginosus sasaimae</i>	N Colombia (upper tropical Magdalena Valley)	Sasaima, 74 kilometers west of Bogota, in the Magdalena Valley.
<i>Clibanornis rubiginosus caquetae</i>	SE Colombia (Caquetá and Putumayo)	Morelia, 900 feet, Caqueta, Colombia.
<i>Thripadectes flammulatus bricenoi</i>	Andes of w Venezuela (Mérida)	Culatd, Andes of Merida, Venezuela.
<i>Thripadectes virgaticeps sclateri</i>	W Andes of Colombia (Cauca and Nariño)	San Pablo, 4500 feet [Province of Tuquerres, Department of Narifio], Colombia.
<i>Thripadectes melanorhynchus striaticeps</i>	E Andes of Colombia (single record from w Meta)	Bogota.
<i>Automolus subulatus nicaraguae</i>	Caribbean lowlands of e Nicaragua	Rio Grande, Nicaragua.
<i>Automolus subulatus cordobae</i>	NW Colombia	Cerro Mucurucucú, Córdoba, Colombia.
<i>Premnoplex brunnescens albescens</i>	Subtropical mountains of e Panama (Cerro Tacarcuna)	east slope of Mt. Tacarcuna, 4600 feet, eastern Panami.

<i>Premnoplex brunnescens coloratus</i>	Subtropical Santa Marta Mountains (ne Colombia)	San Miguel, 7500 feet, Sierra Nevada de Santa Marta, Colombia.
<i>Aphrastura spinicauda bullocki</i>	Mocha I. (Chile)	Mocha Island, Chile.
<i>Aphrastura spinicauda fulva</i>	Chiloé I. (Chile)	Ancud, Chiloe Island, Chile.
<i>Leptasthenura aegithaloides grisescens</i>	Coastal arid s Peru and n Chile	Gatico, Antofagasta, Chile.
<i>Leptasthenura pileata latistriata</i>	Andes of w Peru (Ayacucho)	Manzanallo, cerca de Ticrapo, 13°23'S, 75° 26'W, Castrovirreyna, Huancavelica, Perú.
<i>Leptasthenura andicola extima</i>	Santa Marta Mountains (ne Colombia)	Paramo de Macotama [= Chiruqua], Sierra Nevada de Santa Marta, Colombia.
<i>Leptasthenura andicola exterior</i>	E Andes of Colombia (Boyacá)	Lagunillas, Boyaca, Colombia.
<i>Leptasthenura andicola certhia</i>	Andes of w Venezuela (Mérida and Trujillo)	San Antonio and Paramo de Santo Domingo [Andes of Merida], Venezuela.
<i>Hellmayrea gularis brunneidorsalis</i>	Sierra de Perijá (Colombia/Venezuela border)	Cerro Tetarí, alto Río Negro, Sierra de Perijá, Zulia, Venezuela.
<i>Hellmayrea gularis cinereiventris</i>	Andes of w Venezuela (Mérida and Trujillo)	Quintero, 9250 feet, near Merida, Venezuela.
<i>Asthenes flammulata multostriata</i>	E Andes of Colombia	Bogota.
<i>Asthenes flammulata quindiana</i>	C Andes of Colombia (Nevado de Tolima)	Paramo de Santa Isabel, 12,500 feet, Central Andes of Colombia.
<i>Asthenes wyatti wyatti</i>	Arid Andes of n Colombia (Santander)	Paramo de Pamplona, Santander, Colombia.
<i>Asthenes wyatti sanctaemartae</i>	Santa Marta Mountains (ne Colombia)	Macotama, Sierra Nevada de Santa Marta, Colombia.
<i>Asthenes wyatti mucuchiesi</i>	Andes of nw Venezuela (Mérida)	Paramo de Mucuchies, Merida, Venezuela.
<i>Asthenes wyatti phelpsi</i>	Sierra de Perijá (Colombia/Venezuela border)	"Frontera 2", Sierra de Perijá, Zulia, Venezuela, al pie del Cerro Tetarí, cerca de la frontera con Colombia, 10°58'38'N., 72°35'O.
<i>Asthenes wyatti aequatorialis</i>	W Andes of c Ecuador	Mt. Chimborazo, 13,000 feet, Ecuador.
<i>Asthenes wyatti azuay</i>	Andes of s Ecuador (Azuay)	Bestion, 10,100 feet, Azuay, Ecuador.
<i>Asthenes sclateri cuchacancae</i>	Andes of sw Bolivia (Cochabamba and Potosí)	Cuchacancha, Bolivia.
<i>Asthenes humilis cajamarcae</i>	Arid temperate w Andes of Peru (Cajamarca)	Cajamarca, 10,000 feet, Peru.
<i>Asthenes pyrrholeuca pyrrholeuca</i>	NW Argentina	Paraguay.
<i>Asthenes fuliginosa fumigata</i>	C and Western Andes of Colombia	Páramo del Tábano, Nariño, Colombia.
<i>Xenerpestes minlosi umbraticus</i>	Pacific lowlands of Colombia and northwestern Ecuador	Punto Muchimbo, Río San Juan, El Valle, Colombia.
<i>Roraimia adusta duidae</i>	Tepuis of s Venezuela (Mt. Duida)	Mt. Duida, 5000 feet, Venezuela.
<i>Cranioleuca vulpina apurensis</i>	W Venezuela (w Apure)	Guasualito, 60 meters, upper Apure River, Apure, Venezuela.
<i>Cranioleuca vulpina foxi</i>	C Bolivia (e Cochabamba)	mouth of the Rio Chapare, 825 feet, Cochabamba, Bolivia.
<i>Cranioleuca pyrrhophia rufipennis</i>	Known from type locality in La Paz, Bolivia	Tilotilo, Bolivia.
<i>Cranioleuca semicinerea goyana</i>	E Brazil (s-c Goiás)	Rio das Almas, Fazenda Thome Pinto, 2Vz leagues from Jaragua, Goiaz, Brazil.
<i>Pseudasthenes humicola goodalli</i>	Andes of n Chile (sw Antofagasta)	Quebrada Paposo; Provincia de Antofagasta, Chile.
<i>Pseudasthenes cactorum lachayensis</i>	C coast of Peru (Lomas de Lachay)	Lomas de Lachay, 11°24'S, 77°24'W, Chancay, 90 Km.al norte de Lima, Perú.
<i>Certhiaxis cinnamomeus marabinus</i>	NE Colombia and nw Venezuela	Mene Grande, 80 meters, Lake Maracaibo, Zulia, Venezuela.
<i>Synallaxis gujanensis columbiana</i>	Tropical and subtropical Eastern Andes of Colombia	Buenavista, 4500 feet, Colombia.
<i>Synallaxis brachyura brachyura</i>	Magdalena Valley and n Western Andes of Colombia	Colombia = Bogota.
<i>Synallaxis moesta moesta</i>	Tropical and subtropical Eastern Andes of Colombia	Bogota.
<i>Synallaxis cabanisi fulviventris</i>	E base of Andes of Bolivia (south to s Beni and Cochabamba)	Yungas, 3600 feet, Cochabamba, Bolivia.

<i>Synallaxis albigularis rodolphei</i>	S Colombia	Río Rumiayaco, sureste de Colombia.
<i>Synallaxis albescens latitabunda</i>	SW Costa Rica	Boruca, Costa Rica.
<i>Synallaxis albescens insignis</i>	Andes of Colombia	Quetame, 4800 feet, Eastern Andes, Colombia.
<i>Synallaxis albescens nesiotis</i>	Santa Marta Mts. (ne Colombia) to n Venezuela; Isla Margarita	El Valle, Margarita Island.
<i>Synallaxis azarae media</i>	W and C Andes of Colombia and n Ecuador	Salento, 7000 feet, Central Andes, Colombia.
<i>Synallaxis candei atrigularis</i>	N Colombia (Magdalena Valley)	Gamarra, Magdalena, Colombia.
<i>Synallaxis cinnamomea terrestris</i>	Tobago	Tobago.
<i>Masius chrysopterus bellus</i>	W slope of W Andes to w side of Central Andes of Colombia	Riolima, 4,000 ft., Valle del Cauca, Colombia.
<i>Cryptopipo litae suffusa</i>	Extreme e Panama (Darién) and adjacent nw Colombia	Obaldia, Caribbean slope, eastern Panama.
<i>Lepidothrix coronata caquetae</i>	Tropical se Colombia (east of Andes in w Meta and w Caquetá)	Morelia, Caqueta, Colombia.
<i>Manacus vitellinus milleri</i>	N Colombia in Sinú and lower Cauca valleys	Puerto Valdivia, 360 ft., lower Cauca River, Antioquia, Colombia.
<i>Manacus vitellinus viridiventris</i>	W Colombia w of W Andes and e side in upper Cauca Valley	Jimenez [= Jimenez], near Buenaventura, Pacific slope, Valle del Cauca.
<i>Manacus manacus abditivus</i>	Santa Marta, lower Cauca and middle Magdalena valleys	Sierra Nevada de Santa Marta, Colombia.
<i>Manacus manacus flaveolus</i>	N Colombia (tropical upper Magdalena Valley)	Bogota, Colombia; restricted to Honda, Magdalena Valley, Tolima, Colombia (D. W. S.).
<i>Manacus manacus umbrosus</i>	Tropical s Venezuela (Amazonas)	Cerro Yapacana, upper Orinoco, Amazonas, Venezuela.
<i>Machaeropterus striolatus obscuristriatus</i>	Tropical nw Venezuela (Mérida)	El Vigia, 200 m., Merida, Venezuela.
<i>Pseudopipra pipra minima</i>	W slope of Western Andes of Colombia (Cauca)	Cocal, Andes al oeste de Popayán, Cauca, Colombia (Nuevo nombre para Pipra leucocilla minor pre-ocupado por Pipra mentalis minor Hartert, Chapman, 1914: 623, 1898: 489.
<i>Pseudopipra pipra bolivari</i>	NW Colombia (upper Sinú Valley)	Murucucii, 1,500 m., Bolivar, Colombia
<i>Pseudopipra pipra unica</i>	Subtropical Colombia (Antioquia and Huila)	El Isno [= Lomas de Isnos], head of Magdalena Valley, Huila, Colombia
<i>Ceratopipra erythrocephala flammiceps</i>	E Colombia (Santander)	El Tambor, Santander, Colombia.
<i>Pipreola riefferii tallmanorum</i>	Andes of central Peru (Huánuco)	Hacienda Paty, Cordillera de Carpish, Huánuco, Perú.
<i>Pipreola aureopectus decora</i>	Santa Marta Mountains (ne Colombia)	Chirua, 7,000 ft., Santa Marta region, Colombia.
<i>Pipreola aureopectus festiva</i>	Coastal mountains of n Venezuela	Cumbre de Valencia, Carabobo, Venezuela.
<i>Pipreola formosa rubidior</i>	Mts. of ne Venezuela (Anzoátegui, Monagas and Sucre)	La Trinidad, 5,600 ft., near Carapas, Mount Turumiquire, Sucre, Venezuela.
<i>Phibalura flavirostris boliviana</i>	Foothills of w Bolivia (La Paz)	near Atten [= Aten], Aplobamba [= Apolo], La Paz, Bolivia.
<i>Ampelion rufaxilla antioquiae</i>	Subtropical Andes of w Colombia and (?) n Ecuador	Santa Elena, Antioquia, Colombia.
<i>Pyroderus scutatus occidentalis</i>	Andes of w Colombia and nw Ecuador	San Antonio, 6,600 ft., Western Andes, Colombia.
<i>Pyroderus scutatus granadensis</i>	E Andes of Colombia and w Venezuela	Nouvelle Grenade [= 'Bogota,' Colombia].
<i>Tityra inquisitor buckleyi</i>	Tropical se Colombia (Caquetá) and e Ecuador (Napo-Pastaza)	Yanayacu, eastern Ecuador.
<i>Tityra semifasciata hannumi</i>	Arid tropical nw Mexico (se Sonora and ne Sinaloa)	San Francisco Canon on Sonora-Sinaloa boundary, 2,000 ft., extreme southeastern Sonora, Mexico.
<i>Schiffornis stenorhyncha stenorhyncha</i>	Tropical ne Colombia and n Venezuela	San Esteban, Carabobo, Venezuela.
<i>Laniocera rufescens griseigula</i>	Tropical n Colombia (Córdoba, Antioquia and Santander)	Quimari, 400 m., southwestern Bolivar, Colombia.

<i>Iodopleura pipra leucopygia</i>	Coastal ne Brazil (Paraíba, Pernambuco, Alagoas and Bahia)	British Guiana.
<i>Laniisoma elegans venezuelense</i>	Locally from tropical ne Colombia to nw Venezuela	Santa Barbara, 200 m., Barinas, Venezuela.
<i>Laniisoma elegans cadwaladeri</i>	Tropical nw Bolivia (La Paz)	Santa Ana, 2,200 ft., Rio Coroico, La Paz, Bolivia.
<i>Xenopsaris albinucha minor</i>	Locally in w and c Venezuela and extreme n Brazil (Roraima)	Quiribana de Caicara, Rio Orinoco, Venezuela.
<i>Pachyramphus cinnamomeus magdalenae</i>	N Colombia to nw Venezuela (Maracaibo basin)	Algodonal, Rio Magdalena, Bolivar, Colombia.
<i>Pachyramphus albogriseus coronatus</i>	Santa Marta Mts. (n Colombia) and nw Venezuela (Zulia)	Cerro Tamuypejocha, 1,975 m., upper Rio Negro, Sierra de Perija, Zulia, Venezuela.
<i>Pachyramphus major itzensis</i>	SE Mexico (Campeche, Yucatán, Quintana Roo) and Belize	Chichen Itza, Yucatan, Mexico.
<i>Pachyramphus homochrous canescens</i>	Caribbean coast of n Colombia and nw Venezuela	Cagualito [= Cacagualito] , 1,500 ft.. Sierra Nevada de Santa Marta, Colombia,
<i>Pachyramphus homochrous quimarinus</i>	NW Colombia (Sinú Valley)	Quimari, 400 m., southwestern Bolivar, Colombia.
<i>Oxyruncus cristatus frater</i>	Discontinuously distributed Costa Rica and w Panama	Calovevora, Veraguas, Panama.
<i>Oxyruncus cristatus tocantinsi</i>	Central Brazil (Goiás to Pará and Amapá)	Baiao, Pedral, Rio Tocantins, Brazil.
<i>Onychorhynchus coronatus fraterculus</i>	NE Colombia to e Venezuela (w Zulia and w Barinas)	Sierra Nevada de Santa Marta, Colombia.
<i>Myiobius barbatus semiflavus</i>	E-central Colombia (Nechí region of Antioquia)	El Tambor, Santander, Colombia.
<i>Piprites chloris perijana</i>	Sierra de Perijá (e Colombia) and Andes of w Venezuela	La Sabana, 1,300 m., Rio Negro, Perija, Zulia, Venezuela.
<i>Platyrinchus mystaceus perijanus</i>	Subtropical Sierra de Perijá (Colombia/Venezuela border)	Cerro Pejochaina, 1,800 m., upper Rio Negro, Sierra de Perija, Zulia, Venezuela.
<i>Platyrinchus mystaceus imatacae</i>	S Venezuela (Sierra de Imataca in Bolívar)	Cerro Tomasote, 500 m., El Palmar, Sierra de Imataca, Bolivar, Venezuela.
<i>Platyrinchus mystaceus ptaritepui</i>	Tepuis of se Venezuela (se Bolívar)	Ptari-tepui (southwestern slope), 1,700 m.. Gran Sabana, Bolivar, Venezuela.
<i>Leptopogon amaurocephalus idius</i>	Coiba I. (Panama)	Isla Coiba, Panama.
<i>Leptopogon amaurocephalus diversus</i>	Santa Marta and Magdalena valleys of n Colombia and w Zulia	Mamatoco, Sierra Nevada de Santa Marta, Colombia.
<i>Phylloscartes superciliaris palloris</i>	E Panama (Darién) and adjacent Colombia (Chocó)	Tapalisa [= Cerro Tacarcuna] , Darien, eastern Panama.
<i>Phylloscartes flaviventris</i>	Coastal mts. of n Venezuela	Ejido and Merida, Venezuela; type from Merida, fide Hellmayr, 1927, Field Mus. Nat. Hist., Zool. Ser., 13, pt. 5, p. 349.
<i>Pseudotriccus pelzelni berlepschi</i>	Mountains of extreme e Panama and adjacent nw Colombia	Cerro Pirre, Panama.
<i>Atalotriccus pilaris pilaris</i>	N Colombia and adjacent Venezuela (Zulia and Táchira)	environs of Cartagena, Bolivar, Colombia.
<i>Hemitriccus margaritaceiventer septentrionalis</i>	S Colombia (arid tropical upper Magdalena Valley)	Honda, Magdalena Valley, Colombia.
<i>Hemitriccus margaritaceiventer duidae</i>	Tepuis of s Venezuela (Mt. Duida)	Chorrera de Vegas, 4,700 ft., Cerro Duida, Venezuela.
<i>Hemitriccus margaritaceiventer breweri</i>	Subtropical se Venezuela (Jaua massif)	Meseta de Sarisarihama, 1,400 m., Bolivar, Venezuela,
<i>Hemitriccus granadensis lehmanni</i>	Santa Marta Mountains (ne Colombia)	San Miguel, headwaters of the Rio Macotama, north slope of Sierra Nevada de Santa Marta, Colombia.
<i>Hemitriccus granadensis granadensis</i>	Andes of Colombia and ne Ecuador	Nouvelle-Grenade' [= 'Bogota], Colombia; restricted to Santa Elena, Antioquia, Colombia (J. T. Z.).
<i>Hemitriccus granadensis intensus</i>	Sierra de Perijá (Colombia/Venezuela border)	Cerro Tamuyejocha, 1,975 m, Rio Negro, Sierra de Perija, western Zulia, Venezuela.
<i>Hemitriccus granadensis federalis</i>	Montane forests of coastal n Venezuela (Distrito Federal)	No [= No] Leon, 2,000 m., Distrito Federal, Venezuela.
<i>Hemitriccus granadensis caesius</i>	Andes of se Peru (Puno) and adjacent n Bolivia	Santo Domingo (Inca Mine), 6,000 [ft.] , Puno, southeastern Peru.

<i>Poecilotriccus plumbeiceps viridiceps</i>	S Bolivia to nw Argentina (Jujuy and Salta)	San Lorenzo, Jujuy, Argentina.
<i>Poecilotriccus plumbeiceps cinereipectus</i>	SE Brazil (Espírito Santo and se Minas Gerais); Alagoas	Jatiboca, 900-1,000 m., Espírito Santo, Brazil.
<i>Poecilotriccus latirostris austroriparius</i>	E Brazil (right bank of Rio Tapajós near Santarém in w Pará)	Santarem, Brazil.
<i>Poecilotriccus latirostris senectus</i>	Amazonian Brazil (ne Amazonas and nw Pará)	Boca do Igarape-Piaba, near Obidos, Para, Brazil.
<i>Poecilotriccus sylvia superciliaris</i>	Tropical n Colombia (Cauca and Magdalena valleys)	Venezuela?'; type is labeled Carthagena [= Cartagena, Bolivar, Colombia], fide Hellmayr, 1927, Field Mus. Nat. Hist., Zool. Ser., 13, pt. 5, p. 306.
<i>Rhynchocyclus olivaceus mirus</i>	NW Colombia (lower Atrato Valley and inland from coast)	Murindo, Antioquia, Colombia.
<i>Rhynchocyclus olivaceus flavus</i>	N and c Colombia and n Venezuela	Onaca, 2,500 ft., near coast. Sierra Nevada de Santa Marta, Colombia.
<i>Rhynchocyclus olivaceus tamborensis</i>	Central Colombia (Rio Lebrija region of Santander)	El Tambor, Santander, Colombia.
<i>Tolmomyias sulphurescens asemus</i>	W Colombia (Cauca and Magdalena valleys)	Pavas, western Colombia.
<i>Tolmomyias sulphurescens berlepschi</i>	Trinidad	Caparo, Trinidad.
<i>Tolmomyias sulphurescens duidae</i>	Tepuis of se Venezuela and adjacent nw Brazil	Cano Seco, 2,250 ft., Cerro Duida, Venezuela.
<i>Pyrrhomyias cinnamomeus assimilis</i>	Santa Marta Mountains (ne Colombia)	Valparaiso, Sierra Nevada de Santa Marta, Colombia.
<i>Pyrrhomyias cinnamomeus vieillotioides</i>	Coastal mountains of nw Venezuela (Lara to Miranda)	Caracas, Venezuela.
<i>Pyrrhomyias cinnamomeus spadix</i>	Coastal mountains of ne Venezuela (Anzoátegui to w Sucre)	Los Palmales, 450 m., Monagas, Venezuela.
<i>Camptostoma obsoletum cauae</i>	Colombia (Western Andes, Cauca and Magdalena valleys)	Rio Frio, 3,500 ft., Cauca Valley, Colombia.
<i>Mecocerculus leucophrys montensis</i>	northern Colombia (Sierra Nevada de Santa Marta)	Paramo de Macotama, 11,000 ft., Sierra Nevada de Santa Marta, Colombia.
<i>Mecocerculus leucophrys notatus</i>	W and central Andes of Colombia (south to Cauca)	Leonera, near Caldas, Colombia.
<i>Anairetes alpinus bolivianus</i>	Yungas of w Bolivia (La Paz)	Kilometer 50, 13,500 ft., Yungas railway, La Paz, Bolivia.
<i>Uromyias agraphia plengei</i>	E Peru (Cordillera de Colán in Amazonas)	Cordillera Colán, 5°34'S, 78°19'W, Amazonas, Perú.
<i>Capsiempis flaveola leucophrys</i>	Colombia (Magdalena Valley) to nw Venezuela	Bogota, Colombia.
<i>Capsiempis flaveola magnirostris</i>	SW Ecuador (Pichincha to El Oro)	Chimbo, Ecuador.
<i>Polystictus pectoralis bogotensis</i>	W Colombia (Cauca Valley and Cundinamarca)	Suba, Savannah de Bogota, Colombia.
<i>Myiopagis gaimardii trinitatis</i>	Trinidad	Caparo, Trinidad.
<i>Myiopagis gaimardii bogotensis</i>	Tropical ne Colombia and n Venezuela	Bogota, Colombia.
<i>Myiopagis parambae absita</i>	Extreme e Panama (Darién)	old Tacarcuna village site, 950 m., headwaters of the Rio Pucro, base of Cerro Mali, Serrania del Darien, Panama.
<i>Myiopagis olallai incognita</i>	Serranía de Perijá, Venezuela	Ayapa (=La Sabana), upper Río Negro, State of Zulia, Venezuela (ca. 10°02'N, 72°45'W; 1100 m elevation);
<i>Myiopagis cotta</i>	Jamaica	Jamaica.
<i>Myiopagis viridicata pallens</i>	Colombia (Cundinamarca, Huila and Santa Marta region)	Sierra Nevada de Santa Marta, Colombia.
<i>Elaenia dayi tyleri</i>	tepuis of southern Venezuela (Duida, Huachamacari, and Parú)	Cerro Duida, 5,000 ft., Venezuela.
<i>Elaenia dayi dayi</i>	Tepuis of s Venezuela (Roraima, Kukenamk and Ptari-tepui)	summit of Cerro Roraima, 8,600 ft., Venezuela.
<i>Elaenia frantzii browni</i>	N Colombia (Santa Marta Mountains and Sierra de Perijá)	Pueblo Viejo, 8,000 ft., Sierra Nevada de Santa Marta, Colombia.
<i>Elaenia martinica barbadensis</i>	Barbados	Barbados.

<i>Elaenia martinica chinchorrensis</i>	Great Cay I., Chinchorro Bank, off Quintana Roo (eastern Mexico) and Half Moon Caye, Belize; ; the subspecific identity of elaenias on other cayes off of Belize is not established, and the validity of the poorly known chinchorrensis is questioned by some	Great Key, Chinchorro Bank, Quintana Roo, Mexico.
<i>Elaenia martinica cinerascens</i>	San Andrés, Providencia, Santa Catalina Is. (off Honduras)	Old Providence Island.
<i>Phyllomyias zeledoni zeledoni</i>	Mountains of Costa Rica and w Panama	Dota and Barranca, Costa Rica; restricted to Dota (J. T. Z.).
<i>Phyllomyias zeledoni wetmorei</i>	Sierra de Perijá (Colombia/Venezuela border)	Jamayaujaina [= Cerro Pejochaina] , 1,640 m., Rio Negro, Sierra de Perija, Zulia, Venezuela.
<i>Phyllomyias zeledoni bunites</i>	Tepuis of s Venezuela in Bolívar (Cerro Chimantá-tepui)	western slope of Cerro Chimantatepui, 1,300 m., Bolivar, Venezuela.
<i>Phyllomyias urichi</i>	Extreme ne Venezuela (Sucre, Monagas and Anzoátegui)	Quebrada Seca, Sucre, Venezuela.
<i>Phyllomyias griseiceps caucae</i>	west central Colombia (upper Cauca Valley)	Miraflores, 6,400 ft., east of Palmira, Central Andes, Cauca, Colombia.
<i>Phyllomyias nigrocapillus flavimentum</i>	Santa Marta Mountains (ne Colombia)	San Lorenzo, 7,500 ft. [7,000 ft. on original label] , Sierra Nevada de Santa Marta, Colombia.
<i>Zimmerius chrysops minimus</i>	Santa Marta Mountains (ne Colombia)	Minca, 2,000 ft.. Sierra Nevada de Santa Marta, Colombia.
<i>Euscarthmus meloryphus paulus</i>	NE Colombia (Santa Marta region) and n Venezuela	Chirua, Sierra Nevada de Santa Marta, Colombia.
<i>Lathrotriccus euleri flaviventris</i>	Grenada (Lesser Antilles). Extinct	South America; restricted to Trinidad by Hellmayr, 1927, Field Mus. Nat. Hist., Zool. Ser., 13, pt. 5, p. 215.
<i>Contopus fumigatus cineraceus</i>	Subtropical n Venezuela (Yaracuy to Miranda)	Caracas, Venezuela.
<i>Contopus sordidulus peninsulae</i>	S Baja California; winters to nw South America	Sierra de la Laguna, Baja California, Mexico.
<i>Contopus caribaeus bahamensis</i>	Grand Bahama, Abaco, Andros, New Providence and Eleuthera	Bahamas; cotypes from Nassau, New Providence, fide Bangs and Penard, 1925, Bull. Mus. Comp. Zool., 67, p. 203.
<i>Contopus caribaeus nerlyi</i>	Islands off s Camagüey (Cuba)	Archipelago de los Jardines de la Reina, Camagiiey, Cuba.
<i>Contopus hispaniolensis tacitus</i>	Gonâve I. (Haiti)	Anse a Galets, Gonave Island, Haiti.
<i>Contopus latirostris brunneicapillus</i>	Montane forests of Dominica, Guadeloupe and Martinique	Dominica.
<i>Contopus latirostris latirostris</i>	Montane forests of St. Lucia	St. Lucia.
<i>Empidonax albigularis timidus</i>	Highlands of nw Mexico (Chihuahua to s Durango)	El Salto, Durango, Mexico.
<i>Empidonax affinis vigensis</i>	Oak-pine forests of e Mexico (Veracruz)	Veracruz, Mexico.
<i>Empidonax fulvifrons fulvifrons</i>	Mountains of ne Mexico (Tamaulipas and San Luis Potosí)	Texas; ?error, probably mountains of northeastern Mexico as suggested by Ridgway, 1884, Proc. Biol. Soc. Washington, 2, p. 109; restricted to Miquiahuana, Tamaulipas by Lowery and Dalquest, 1951, Univ. Kansas Publ., Mus. Nat. Hist., 3, p. 612.
<i>Empidonax fulvifrons brodkorbi</i>	S Mexico (Río Molino area of s Oaxaca)	Río Molino (ca. lat. 16° 5' N., long. 96° 29' W.), southern Oaxaca, Mexico.
<i>Empidonax fulvifrons inexpectatus</i>	Central and s Honduras	Cerro Cantoral, 6,500 ft., Achaga, Honduras.
<i>Sayornis nigricans aquaticus</i>	S Mexico (highlands of s Chiapas) to Guatemala and Nicaragua	Duehas, Guatemala.
<i>Sayornis nigricans amnicola</i>	Highlands of Costa Rica and w Panama (Chiriquí)	Boquete, Panama.
<i>Sayornis saya quiescens</i>	Northern half of Baja California and Isla de Cedros	San Jose, 2,500 ft., about lat. 31° N., about 45 miles northeast of San Quintin, Baja California, Mexico.
<i>Sublegatus arenarum tortugensis</i>	Isla La Tortuga (off n Venezuela)	Garambeo, Isla La Tortuga, Venezuela.
<i>Pyrocephalus rubinus major</i>	SE Peru (Cuzco and Puno)	no locality; restricted to Chaquimayo, Andes of Carabaya, Peru by Laubmann, 1930, Deutsch. Gran Chaco-Exped. Vogel, p. 214.

<i>Pyrocephalus nanus nanus</i>	Galapagos Islands (except San Cristóbal)	Galapagos Archipelago; type from James Island [= San Salvador] , fide Ridgway, 1894, Proc. U. S. Nat. Mus., 17, p. 365
<i>Pyrocephalus nanus dubius</i>	extinct; formerly San Cristóbal I. (Galapagos Islands); last reported 1987	Galapagos (Colon) Islands [= Chatham Island (Isla San Cristobal)]
<i>Knipolegus poecilurus venezuelanus</i>	Coastal cordillera of n Venezuela (Distrito Federal)	El Escorial, 2,500 m., Merida, Venezuela.
<i>Muscisaxicola maculirostris niceforoi</i>	Eastern Andes of Colombia (Boyacá and Cundinamarca)	mountains of Bogota, Colombia.
<i>Muscisaxicola maculirostris rufescens</i>	Andes of Ecuador (Pichincha to Azuay)	Yocon, Chimborazo, Ecuador.
<i>Muscisaxicola rufivertex rufivertex</i>	Andes of s Chile and s Argentina; winters north to Antofagasta	Cobija, Antofagasta, Chile.
<i>Muscisaxicola maclovianus maclovianus</i>	Falkland Islands	lies Malouines, dans les environs de la baie de l'Huile, du port Duperry et les vallees qui avoisinent l'Aiguade.
<i>Muscisaxicola alpinus columbianus</i>	Central Andes of Colombia (Nevado del Ruiz to Cauca)	Paramo de Santa Isabel, Central Andes, Colombia.
<i>Muscisaxicola alpinus quesadae</i>	E Andes of Colombia (Boyacá and Cundinamarca)	Lagunillas, Boyaca, Colombia.
<i>Cnemarchus erythropygius orinomus</i>	Santa Marta Mountains and Eastern Andes of n Colombia	between 11,400 and 12,000 ft., above Mamancanaca, Sierra Nevada de Santa Marta, Colombia.
<i>Pyrope pyrope fortis</i>	Chiloé I. (Chile)	Ancud, Isla Grande de Chiloé, Chile
<i>Agriornis albicauda pollens</i>	Andes of Ecuador (north to Imbabura)	New name for <i>Agriornis andicola</i> Sclater PZS 1860, p.78; Panza, Ecuador.
<i>Agriornis lividus lividus</i>	Coast and mountains of Chile (Atacama to Valdivia)	mountains of Valparaíso, Chile.
<i>Silvicultrix diadema towarensis</i>	Coastal cordillera of n Venezuela (Aragua and Distrito Federal)	Colonia Tovar, 1,950 m., Aragua, Venezuela.
<i>Ochthoeca rufipectoralis poliogastra</i>	Santa Marta Mountains (ne Colombia)	Sierra Nevada de Santa Marta, Colombia.
<i>Ochthoeca rufipectoralis rubicundula</i>	Sierra de Perijá (Colombia/Venezuela border)	above Airoca, between 9,500 and 10,000 ft.. Sierra de Perija, Magdalena, Colombia.
<i>Ochthoeca rufipectoralis rufopectus</i>	Eastern Andes of Colombia (south to Bogotá)	Colombia; restricted to Pacho, Cundinamarca by Zimmer, 1942, Amer. Mus. Novit., no 1203, p. 14.
<i>Ochthoeca fumicolor ferruginea</i>	Central and Western Andes of Colombia (Antioquia)	Santa Elena, 9,000 ft., Antioquia, Colombia.
<i>Ochthoeca leucophrys leucophrys</i>	Andes of w Bolivia	Sicasica, La Paz, Bolivia.
<i>Ramphotrigon megacephalum venezuelense</i>	NW Venezuela (and probably adjacent Colombia)	Ciudad Bolivia, 150 m., Barinas, Venezuela.
<i>Attila spadiceus caniceps</i>	Trop. n Colombia (middle Magdalena and lower Sinú valleys)	Jaraquiel, Bolívar [= Córdoba], Colombia.
<i>Attila spadiceus parvirostris</i>	Santa Marta Mountains (ne Colombia) and nw Venezuela	Minca, Sierra Nevada de Santa Marta, Colombia.
<i>Myiarchus yucatanensis lanyoni</i>	Cozumel I. (off Yucatán Peninsula of e Mexico)	ca. 12 km. southwest of San Miguel, Isla Cozumel, Quintana Roo, Mexico.
<i>Myiarchus cephalotes caribbaeus</i>	Mountains of n Venezuela (Trujillo and Lara to Sucre)	Galipan, 6,000 ft., Cerro del Avila, Distrito Federal, Venezuela.
<i>Myiarchus tyrannulus cozumelae</i>	breeds Cozumel I. (off Yucatán Peninsula of southeastern Mexico); winter range unknown	Cedral, Isla Cozumel, Quintana Roo, México
<i>Myiarchus tyrannulus insularum</i>	Utila, Roatán and Bonaca islands (off Honduras)	Isla Utila, Honduras.
<i>Myiarchus stolidus stolidus</i>	Jamaica	Jamaica.
<i>Myiarchus oberi oberi</i>	Lesser Antilles (Dominica and Guadeloupe)	Dominica.
<i>Myiarchus oberi sanctaeluciae</i>	St. Lucia (Lesser Antilles)	Santa Lucia.
<i>Myiarchus oberi sclateri</i>	Martinique (Lesser Antilles)	Martinique.
<i>Machetornis rixosa obscuradorsalis</i>	llanos of eastern Colombia and southwestern Venezuela; records from northeastern Ecuador presumably are of this subspecies	Ciudad Bolivia, 150 m., Barinas, Venezuela.

<i>Pitangus sulphuratus caucensis</i>	W and s Colombia (sw Bolívar, Cauca and Magdalena valleys)	Cali, Cauca Valley, Colombia.
<i>Megarynchus pitangua tardiusculus</i>	Foothills of nw Mexico (s Sinaloa to Nayarit)	Rancho Santa Barbara, 2,500 ft., 20 miles northeast of Rosario, Sinaloa, Mexico.
<i>Megarynchus pitangua caniceps</i>	SW Mexico (s Jalisco)	Barranca Veltran, near San Marcos, southern Jalisco, Mexico.
<i>Megarynchus pitangua deserticola</i>	Central Guatemala (valley of Río Negro)	Sacapulas, 25 miles east of Nebaj, Río Negro valley, Guatemala.
<i>Myiozetetes cayanensis rufipennis</i>	E Colombia to n Venezuela and e Ecuador	Valencia, Carabobo, Venezuela.
<i>Conopias albobittatus distinctus</i>	Lower Caribbean slopes of Costa Rica	Rio Reventazon, near Guayabo, Costa Rica.
<i>Conopias cinchoneti icterophrys</i>	Andes of Colombia to nw Venezuela (Sierra de Perijá)	Bogota, ' Colombia.
<i>Myiodynastes maculatus nobilis</i>	Caribbean coast of ne Colombia to w slope of Sierra de Perijá	Sierra Nevada de Santa Marta, Colombia.
<i>Tyrannus vociferans xenopterus</i>	Highlands of sw Mexico (Guerrero)	Chilpancingo, Guerrero, Mexico.
<i>Tyrannus dominicensis vorax</i>	Lesser Antilles; winters to Trinidad and the Guianas	Martinique.
<i>Tyrannus caudifasciatus flavescens</i>	Isle of Pines	Los Indios, Isle of Pines.
<i>Tyrannus caudifasciatus caymanensis</i>	Cayman Islands	Grand Cayman, West Indies.
<i>Tyrannus savana sanctaemartae</i>	Caribbean coast of n Colombia and nw Venezuela (Guajira Pen.)	Bonda, Sierra Nevada de Santa Marta, Colombia.